


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

B12

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : A61K 38/00, 38/04, C07K 5/00, 7/00, 16/00	A1	(11) International Publication Number: WO 96/01643 (43) International Publication Date: 25 January 1996 (25.01.96)
(21) International Application Number: PCT/US95/08401 (22) International Filing Date: 30 June 1995 (30.06.95) (30) Priority Data: 08/271,943 8 July 1994 (08.07.94) US (60) Parent Application or Grant (63) Related by Continuation US 08/271,943 (CIP) Filed on 8 July 1994 (08.07.94) (71) Applicants (for all designated States except US): THOMAS JEFFERSON UNIVERSITY [US/US]; 11th and Walnut Streets, Philadelphia, PA 19107 (US). KING'S COLLEGE [GB/GB]; Biomedical Sciences Division The Randall Insti- tute, 29/29 Drury Lane, London WC2B 5R1 (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): JAMESON, Bradford, A. [US/US]; 810 West Sedgewick Street, Philadelphia, PA 19119 (US). SUTTON, Brian, J. [GB/GB]; 16 Playfield Road, Kennington, Oxford OX1 5SR (GB). McDONNELL, James, M. [US/GB]; King's College, Biomedical Sciences Division The Randall Institute, 29/29 Drury Lane, London	WC2B 5R1 (GB). GOULD, Hannah, J. [US/GB]; 48 Great Percy Street, London WC1 (GB). KORNGOLD, Robert [US/US]; 101 Dalton Terrace, Cherry Hill, NJ 08003 (US). BEAVIL, Andrew, J. [GB/GB]; 7 Dobson Road, Gravesend, Kent DA12 5TF (GB). (74) Agents: JOHNSON, Philip, S. et al.; Woodcock Washburn Kurtz Mackiewicz & Norris, 46th floor, One Liberty Place, Philadelphia, PA 19103 (US). (81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published With international search report.	
(54) Title: IgE ANTAGONISTS (57) Abstract <p>Peptides and peptide dimers which are capable of modulating IgE binding of FcεR1 are disclosed. Pharmaceutical compositions comprising such peptides and dimers are disclosed. Methods of inhibiting IgE binding to FcεR1 on cells and therapeutically or prophylactically treating an individual suspected of suffering from or being susceptible to an immediate allergic response are disclosed.</p>		



IgE Antagonists

Field of the Invention

The present invention relates to biologically active compounds which are similar to regions of the high affinity Fc
5 réceptor for immunoglobulin E.

Background of the Invention

Clinical bone marrow transplantation is an important therapeutic treatment for several diseases including high risk leukemia, aplastic anemia, and severe combined
10 immunodeficiency. In addition, there is a wide range of metabolic and genetic disorders that can potentially be corrected by this approach. However, the usefulness of marrow transplantation is currently limited by several important risk factors, the principal one being graft-versus-host disease
15 (GVHD), an often times lethal complication which occurs in a high proportion of transplants.

The risk of GVHD can be reduced by HLA matching of the marrow donor and recipient, with a matched sibling being the primary choice. Yet, less than 30% of the patients in North
20 America have an HLA-matched sibling, and therefore must seek suitable unrelated HLA-matched donors from the National Marrow Donor Program. The probability of finding an unrelated HLA-matched donor is currently on the order of 30-40% and depend on the total number of donors registered. In both related and
25 unrelated HLA-matched transplant situations, the risk of GVHD is still quite high due to disparity of non-HLA multiple minor histocompatibility (H) antigens. GVHD is somewhat higher in

- 2 -

unrelated cases, as this increases the probability of differences at these loci.

Mature donor T cells contaminating the marrow inoculum are responsible for GVHD. Several studies have shown that depletion of these T cells significantly diminishes the incidence of disease. However, the elimination of donor T cells has also resulted in a greater incidence of leukemic relapse. It seems important to provide at least some level of T cell immunocompetency in these completely immunocompromised patients to not only combat residual leukemia cells but also to counter opportunistic infections. In this respect, the same GVHD-reactive donor T cells may be important for targeting leukemia cells expressing the same host allogeneic histocompatibility antigens. Therapeutic approaches that could ameliorate the pathogenic tissue destruction accompanying GVHD, particularly in the gut and skin, but that would allow for continued anti-leukemia activity would greatly benefit marrow transplant patients.

Immediate allergic responses, also referred to as type 1 hypersensitivity reactions, are mediated through the interaction of immunoglobulin E (IgE) with the α -chain of its high affinity Fc receptor, Fc ϵ R1. The binding of a multivalent allergen to an IgE-Fc ϵ R1 complex initiates a cross-linking of the receptor and consequent cellular activation. These high affinity receptors are found primarily on mast cells and basophils. When activated by an allergen, these cells respond by releasing histamine, eicosanoids and cytokines.

In addition to its high affinity Fc receptor, IgE binds to a low affinity receptor, the Fc ϵ R2, which is found on B cells, T cells, macrophages, NK cells, eosinophils, platelets, follicular dendritic cells as well as several other cell types. Activation of the Fc ϵ R2 has been implicated in IgE-dependent cell cytotoxicity (ADCC) as well as in allergic inflammation.

Many different fine mapping studies have been conducted on the binding of the IgE to its Fc receptors. The consensus mapping data is confirmed by a recent study by Nissim

- 3 -

et al. (1993) *J. Immunol.* 150:1365-1374, which is incorporated herein by reference, showing that the binding activity for both the high and low affinity receptors resides within the C ϵ 3 (the third constant domain) of the IgE. Data show that the species
5 specific binding of IgE to the Fc ϵ R1 is contained within the first 16 amino acids of the C ϵ 3, whereas no species specific binding is observed in the same region with respect to binding the Fc ϵ R2.

There is a need for compounds and methods which can
10 inhibit IgE-Fc ϵ R1 interaction. IgE-Fc ϵ R1 interaction is associated with immediate allergic reactions. There is a need for compounds and methods which can inhibit mast cell and basophil activity that results from IgE-Fc ϵ R1 interaction. Mast cell and basophil activity linked to IgE-Fc ϵ R1 interaction
15 is associated, for example with reactions to allergens and GVHD. Identification of compounds that inhibit the IgE-Fc ϵ R1 interaction can be used in methods of treating allergies and GVHD.

Summary of the Invention

20 The present invention relates to peptides that modulate IgE binding to Fc ϵ R1 and that consist of 4-50 amino acids and comprise at least a fragment of SEQ ID NO:3 of at least 4 amino acids.

The present invention relates to pharmaceutical
25 compositions comprising a peptide that modulate IgE binding to Fc ϵ R1 and that consist of 4-50 amino acids and comprises at least a fragment of SEQ ID NO:3 of at least 4 amino acids and a pharmaceutically acceptable carrier or diluent.

The present invention relates to peptides that
30 modulate IgE binding to Fc ϵ R1 and that consist of 6-50 amino acids and comprise SEQ ID NO:5.

The present invention relates to pharmaceutical compositions comprising a peptide that modulate IgE binding to Fc ϵ R1 and that consist of 6-50 amino acids and comprise SEQ ID
35 NO:5 and a pharmaceutically acceptable carrier or diluent.

- 4 -

The present invention relates to a method of inhibiting IgE binding to FcεR1 on cells comprising the steps of contacting cells that contain the FcεR1 with a peptide that inhibits IgE binding to FcεR1 and that consists of 4-50 amino acids and comprises at least a fragment of SEQ ID NO:3 of at least 4 amino acids in an amount sufficient to inhibit IgE binding to FcεR1.

The present invention relates to a method of inhibiting IgE binding to FcεR1 on cells comprising the steps of contacting cells that contain the FcεR1 with a peptide that inhibits IgE binding to FcεR1 and that consists of 6-50 amino acids and comprises SEQ ID NO:5 in an amount sufficient to inhibit IgE binding to FcεR1.

The present invention relates to a method of therapeutically or prophylactically treating an individual suffering from or susceptible to an immediate allergic response comprising administering to such an individual an effective amount of a peptide that inhibits IgE binding to FcεR1 and that consists of 4-50 amino acids and comprises at least a fragment of SEQ ID NO:3 of at least 4 amino acids in an amount sufficient to inhibit IgE binding to FcεR1.

The present invention relates to a method of therapeutically or prophylactically treating an individual suffering from or susceptible to an immediate allergic response comprising administering to such an individual an effective amount of a peptide that inhibits IgE binding to FcεR1 and that consists of 6-50 amino acids and comprises SEQ ID NO:5 in an amount sufficient to inhibit IgE binding to FcεR1.

The present invention relates to a peptide dimer that modulate IgE binding to FcεR1 comprising a first monomer which consists of between 4 and 50 amino acids and comprises at least a fragment of SEQ ID NO:3 of at least 4 amino acids, and a second monomer which consists of between 6 and 50 amino acids and comprises SEQ ID NO:5.

35 Brief Description of the Drawings

- 5 -

Figure 1 shows data from *in vitro* experiments testing the effect of various concentrations of D amino acid peptide #1, which is described in Example 3, on the level of inhibition of IgE-IgE receptor.

5 Figure 2 shows data from *in vivo* experiments testing the effect of D amino acid peptide #1, which is described in Example 3, on survival of mice in a GVHD model.

Figure 3 shows data from inhibition assays using a biosensor in which the association of IgE-Fc at a concentration
10 of 5nM was monitored for 6 minutes in the presence at various concentrations and absence of a biotinylated derivative of the D amino acid peptide #1, which is described in Example 3.

Figure 4 shows the *in vitro* inhibition of binding of IgE Fc to Fcε by biosensor assay.

15 Figure 5 shows *in vivo* data that demonstrates inhibition of GVHD by compounds of the invention.

Figure 6 shows *in vivo* data that demonstrates inhibition of antigen-induced anaphylaxis by compounds of the invention.

20 Description of Preferred Embodiments

The present invention provides conformationally restrained analogs which are modelled from the FcεR1 α-chain surface. This application is related to U.S. Patent Application Serial Number 08/271,943 which is incorporated in
25 its entirety herein by reference. Using commercially available molecular modeling software, a structural model of the α-chain of the IgE high affinity receptor was created using the known three-dimensional coordinates of related proteins (See Beavil, A.J. et al. 1993 *Biochem. Soc. Trans.* 21:968-972, which is
30 incorporated herein by reference). CD2 was chosen as the principal template because its three dimensional structure was known and it shares significant sequence homology with the IgE-receptor protein.

The compounds of the invention are capable of
35 inhibiting IgE-IgE receptor binding linked to immediate allergic responses. The compounds of the invention intervene

- 6 -

in the IgE-IgE receptor signal pathway and, thereby, creating therapeutic reagents for the treatment of IgE-dependent disease states including allergic rhinitis, food allergies, atopic dermatitis and allergic asthma.

5 The amino acid sequence of FcεR1 is shown in SEQ ID NO:1. The amino acid sequence of CD2 is shown in SEQ ID NO:2. The known three dimensional structure of CD2 was used as a template. The FcεR1 amino acid sequence shown in SEQ ID NO:1 was aligned to the three dimensional CD2 template. Alignment
10 of the sequences allowed for a prediction of the three dimensional structure of FcεR1. The FcεR1 IgE-binding region in the α-chain sequence was identified using the predicted three dimensional structure. The modeled structure of the α-chain of the high affinity IgE receptor, in turn, was used as
15 a template in selecting and designing peptides, particularly conformationally restrained analogs of the receptor surface.

The location of a principal IgE-binding region in the α-chain of FcεR1 occurs between amino acids 140 and 160 (134-154 of SEQ ID NO:1), more particularly between amino acids 144-
20 154 (139-149 of SEQ ID NO:1) of the FcεR1 protein sequence. The amino acid sequence 144-154 (139-149 of SEQ ID NO:1) is shown as SEQ ID NO:3.

According to some embodiments, the peptides of the invention that inhibit IgE from binding to FcεR1 consist of 4-
25 50 amino acids and comprise at least a fragment of SEQ ID NO:3 of at least 4 amino acids.

In some preferred embodiments, the peptides of the invention consist of 4-25 amino acids including at least a fragment of SEQ ID NO:3 of at least 4 amino acids. In some
30 embodiments, the peptide of the invention consist of 6-20 amino acids including at least a fragment of SEQ ID NO:3 of at least 4 amino acids. In some embodiments, the peptide of the invention consist of 8-15 amino acids including at least a fragment of SEQ ID NO:3 of at least 4 amino acids. In some
35 embodiments, the peptides of the invention consist of 10-12 amino acid residues including at least a fragment of SEQ ID NO:3 of at least 4 amino acids.

- 7 -

As used herein, the term "fragment of SEQ ID NO:3" is meant to refer to peptides which comprise an amino acid sequence identical to a portion of SEQ ID NO:3 having at least 4. In some embodiments, peptides that comprise a fragment of SEQ ID NO:3 include fragments of SEQ ID NO:3 having 4, 5, 6, 7, 8, 9 or 10. Accordingly, a fragment of SEQ ID NO:3 of at least 4 amino acids may be have amino acids 1-4, 2-5, 3-6, 4-7, 5-8, 6-9, 7-10, 8-11, 1-5, 2-6, 3-7, 4-8, 5-9, 6-10, 7-11, 1-6, 2-7, 3-8, 4-9, 5-10, 6-11, 1-7, 2-8, 3-9, 4-10, 5-11, 1-8, 2-9, 3-10, 4-11, 1-9, 2-10, 3-11, 1-10 or 2-11 of SEQ ID NO:3. Peptides may comprise SEQ ID NO:3. Thus, a peptide of the invention that comprises at least a fragment of SEQ ID NO:3 of at least 4 amino acids may comprise SEQ ID NO:3 or a fragment thereof such as one that has amino acids 1-4, 2-5, 3-6, 4-7, 5-8, 6-9, 7-10, 8-11, 1-5, 2-6, 3-7, 4-8, 5-9, 6-10, 7-11, 1-6, 2-7, 3-8, 4-9, 5-10, 6-11, 1-7, 2-8, 3-9, 4-10, 5-11, 1-8, 2-9, 3-10, 4-11, 1-9, 2-10, 3-11, 1-10 and 2-11 of SEQ ID NO:3. In some preferred embodiments, the peptides of the invention comprise SEQ ID NO:3. In some preferred embodiments, the peptides comprise a fragment of SEQ ID NO:3 that consists of 4-10 amino acids. In some embodiments, the peptides comprise a fragment of SEQ ID NO:3 that consists of 6-10 amino acids. In some embodiments, the peptides comprise a fragment of SEQ ID NO:3 that consists of 8-10 amino acids.

In addition to SEQ ID NO:3 or fragments thereof, peptides of the invention may comprise at least a fragment of SEQ ID NO:3 of at least 4 amino acids and further comprise additional amino acids. The additional amino acids may be FcεR1 amino acids, non-FcεR1 amino acids or both. FcεR1 amino acids include amino acids 140-143 (amino acids 134-137 of SEQ ID NO:1) or fragments thereof, amino acids 155-160 (amino acids 149-154 of SEQ ID NO:1) or fragments thereof. Additional FcεR1 amino acids include adjacent amino acid sequences. Moreover, peptides may comprise other FcεR1 amino acid sequences such as those which define other IgE-FcεR1 binding regions. Non-FcεR1 amino acids include amino acids used to conformationally restrict the peptide.

- 8 -

In some preferred embodiments, peptides are conformationally constrained and consist of amino and carboxy terminal cysteines flanking SEQ ID NO:3 or a fragment thereof selected from the group consisting of amino acids 1-4, 2-5, 3-6, 4-7, 5-8, 6-9, 7-10, 8-11, 1-5, 2-6, 3-7, 4-8, 5-9, 6-10, 7-11, 1-6, 2-7, 3-8, 4-9, 5-10, 6-11, 1-7, 2-8, 3-9, 4-10, 5-11, 1-8, 2-9, 3-10, 4-11, 1-9, 2-10, 3-11, 1-10 or 2-11 of SEQ ID NO:3. One or more residues of such peptides may be a D amino acid. Peptides may be biotinylated. The amino acid sequences described herein may be constructed to proceed in sequence from the amino terminus to carboxy terminus or from the carboxy terminus to the amino terminus. When the sequence proceeds from the amino terminus to carboxy terminus, it is usually constructed of L amino acids. In some embodiments, it is constructed of all L amino acids except either an N terminal cysteine or a C terminal cysteine which is a D amino acid. When D amino acids are used to construct the compound, the sequence is usually reversed from the order it occurs as an L amino acid peptide. It therefore proceeds from the carboxy terminus to the amino terminus. In some embodiments, it is constructed of all D amino acids except either an N terminal cysteine or a C terminal cysteine which is an L amino acid.

The location of another IgE-binding region of FcεR1 occurs between amino acids 168-186 of SEQ ID NO:1, more particularly between amino acids 174-179 of SEQ ID NO:1. The amino acid sequence 168-186 of SEQ ID NO:1 is shown as SEQ ID NO:4. The amino acid sequence 174-179 of SEQ ID NO:1 is shown as SEQ ID NO:5.

In some embodiments, the peptides of the invention that inhibit IgE from binding to FcεR1 consist of 6-50 amino acids and comprises SEQ ID NO:5.

In some preferred embodiments, the peptides of the invention consist of 6-25 amino acids including SEQ ID NO:5. In some embodiments, the peptides consist of 6-20 amino acids including SEQ ID NO:5. In some embodiments, the peptides consist of 8-15 amino acids including SEQ ID NO:5. In some

- 9 -

embodiments, the peptides consist of 10-12 amino acids including SEQ ID NO:5.

In some embodiments, peptides of the invention which comprise SEQ ID NO:5 may comprise SEQ ID NO:4 or a fragment of
5 SEQ ID NO:4 of at least 6 amino acids including SEQ ID NO:5. As used herein, the term "fragment of SEQ ID NO:4 of at least 6 amino acids including SEQ ID NO:5" is meant to refer to peptides which comprise an amino acid sequence identical to a portion of SEQ ID NO:4 having at least 6 including amino acids
10 7-12 of SEQ ID NO:4. Amino acids 7-12 of SEQ ID NO:4 are SEQ ID NO:5. In some embodiments, peptides that comprise a fragment of SEQ ID NO:4 having at least 6 amino acids including SEQ ID NO:5 include fragments of SEQ ID NO:4 having 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18 or 19 amino acids.
15 Accordingly, a fragment of SEQ ID NO:4 of at least 6 amino acids including SEQ ID NO:5 may be have amino acids 7-12, 7-13, 7-14, 7-15, 7-16, 7-17, 7-18, 7-19, 6-12, 6-13, 6-14, 6-15, 6-16, 6-17, 6-18, 6-19, 5-12, 5-13, 5-14, 5-15, 5-16, 5-17, 5-18, 5-19, 4-12, 4-13, 4-14, 4-15, 4-16, 4-17, 4-18, 4-19, 3-12, 3-
20 13, 3-14, 3-15, 3-16, 3-17, 3-18, 3-19, 2-12, 2-13, 2-14, 2-15, 2-16, 2-17, 2-18, 2-19, 1-12, 1-13, 1-14, 1-15, 1-16, 1-17, 1-18 or 1-19 of SEQ ID NO:4. Peptides may comprise SEQ ID NO:4. Thus, a peptide of the invention that comprises at least a fragment of SEQ ID NO:4 of at least 6 amino acids including SEQ
25 ID NO:5 may comprise SEQ ID NO:4 or a fragment thereof such as one that has amino acids 7-12, 7-13, 7-14, 7-15, 7-16, 7-17, 7-18, 7-19, 6-12, 6-13, 6-14, 6-15, 6-16, 6-17, 6-18, 6-19, 5-12, 5-13, 5-14, 5-15, 5-16, 5-17, 5-18, 5-19, 4-12, 4-13, 4-14, 4-15, 4-16, 4-17, 4-18, 4-19, 3-12, 3-13, 3-14, 3-15, 3-16, 3-17,
30 3-18, 3-19, 2-12, 2-13, 2-14, 2-15, 2-16, 2-17, 2-18, 2-19, 1-12, 1-13, 1-14, 1-15, 1-16, 1-17, 1-18 and 1-19 of SEQ ID NO:4. In some preferred embodiments, the peptides comprise a fragment of SEQ ID NO:4 that consists of 6-10 amino acids including SEQ ID NO:5. In some embodiments, the peptides comprise a fragment
35 of SEQ ID NO:4 that consists of 6-10 amino acids including SEQ ID NO:5. In some embodiments, the peptides comprise a fragment

- 10 -

of SEQ ID NO:4 that consists of 8-10 amino acids including SEQ ID NO:5.

In addition to SEQ ID NO:5 or fragments of SEQ ID NO:4 that comprise SEQ ID NO:5, according to some embodiments, 5 peptides of the invention may comprise additional amino acids. The additional amino acids may be FcεR1 amino acids, non-FcεR1 amino acids or both. Additional FcεR1 amino acids include adjacent amino acid sequences such as amino acids 150-167 and 187-200 of SEQ ID NO:1 or fragments thereof. Moreover, 10 peptides may comprise other FcεR1 amino acid sequences such as those which define other IgE-FcεR1 binding regions. Non-FcεR1 amino acids include amino acids used to conformationally restrict the peptide.

In some preferred embodiments, peptides are 15 conformationally constrained and consist of amino and carboxy terminal cysteines flanking SEQ ID NO:4 or a fragment thereof selected from the group consisting of amino acids 7-12, 7-13, 7-14, 7-15, 7-16, 7-17, 7-18, 7-19, 6-12, 6-13, 6-14, 6-15, 6-16, 6-17, 6-18, 6-19, 5-12, 5-13, 5-14, 5-15, 5-16, 5-17, 5-18, 20 5-19, 4-12, 4-13, 4-14, 4-15, 4-16, 4-17, 4-18, 4-19, 3-12, 3-13, 3-14, 3-15, 3-16, 3-17, 3-18, 3-19, 2-12, 2-13, 2-14, 2-15, 2-16, 2-17, 2-18, 2-19, 1-12, 1-13, 1-14, 1-15, 1-16, 1-17, 1-18 or 1-19 of SEQ ID NO:4. One or more residues of such peptides may be a D amino acid. Peptides may be biotinylated. 25 The amino acid sequences described herein may be constructed to proceed in sequence from the amino terminus to carboxy terminus or from the carboxy terminus to the amino terminus. When the sequence proceeds from the amino terminus to carboxy terminus, it is usually constructed of L amino acids. In some 30 embodiments, it is constructed of all L amino acids except either an N terminal cysteine or a C terminal cysteine which is a D amino acid. When D amino acids are used to construct the compound, the sequence is usually reversed from the order it occurs as an L amino acid peptide. It therefore proceeds from 35 the carboxy terminus to the amino terminus. In some embodiments, it is constructed of all D amino acids except

- 11 -

either an N terminal cysteine or a C terminal cysteine which is an L amino acid.

According to some embodiments of the invention, peptide dimers are provided which comprise two different monomeric subunits, one of which is a peptide of the invention that comprises at least a fragment of SEQ ID NO:3 of at least 4 amino acids and the other of which is a peptide of the invention which comprises SEQ ID NO:5. Dimeric peptides of the invention inhibit IgE from binding to FcεR1 and comprise one monomer that consists of 4-50 amino acids including at least a fragment of SEQ ID NO:3 of at least 4 amino acids, and one monomer that consists of 6-50 amino acids including SEQ ID NO:5.

Dimers according to the invention comprise two monomeric subunits linked to each other by non-peptide bonds. In some preferred dimers, each of the monomeric subunits has a cysteine residue at at least one terminus and the dimer is formed by disulfide bonds which form between a terminal cysteines of each monomer. In some preferred dimers, the monomeric subunits are linked by other covalent bonds between terminal residues. Methods of linking monomers to form dimers are well known to those having ordinary skill in the art.

In some embodiments, the amino terminal of the monomer that comprises at least a fragment of SEQ ID NO:3 is linked to the amino terminal of the monomer that comprises SEQ ID NO:5. In some embodiments, the amino terminal of the monomer that comprises at least a fragment of SEQ ID NO:3 is linked to the carboxy terminal of the monomer that comprises SEQ ID NO:5. In some embodiments, the carboxy terminal of the monomer that comprises at least a fragment of SEQ ID NO:3 is linked to the amino terminal of the monomer that comprises SEQ ID NO:5. In some embodiments, the carboxy terminal of the monomer that comprises at least a fragment of SEQ ID NO:3 is linked to the carboxy terminal of the monomer that comprises SEQ ID NO:5.

In preferred embodiments of the invention which relate to dimers, the monomer that comprises at least a fragment of SEQ ID NO:3 of at least 4 amino acids consists of 4-25 amino

- 12 -

acids. In some embodiments, the monomer that comprises at least a fragment of SEQ ID NO:3 consists of 6-20 amino acids. In some embodiments, the monomer that comprises at least a fragment of SEQ ID NO:3 consists of 8-15 amino acids. In some
5 embodiments, the monomer that comprises at least a fragment of SEQ ID NO:3 consists of 10-12 amino acid residues.

In preferred embodiments of the invention which relate to dimers, the monomer that comprises at least a fragment of SEQ ID NO:3 of at least 4 amino acids comprises SEQ ID NO:3 or
10 a fragment thereof such as one that has amino acids 1-4, 2-5, 3-6, 4-7, 5-8, 6-9, 7-10, 8-11, 1-5, 2-6, 3-7, 4-8, 5-9, 6-10, 7-11, 1-6, 2-7, 3-8, 4-9, 5-10, 6-11, 1-7, 2-8, 3-9, 4-10, 5-11, 1-8, 2-9, 3-10, 4-11, 1-9, 2-10, 3-11, 1-10 and 2-11 of SEQ ID NO:3.

15 In preferred embodiments of the invention which relate to dimers, the monomer that comprises SEQ ID NO:5 consists of 6-25 amino acids. In some embodiments, the monomer which comprises SEQ ID NO:5 consists of 6-20 amino acids. In some embodiments, the monomer that comprises SEQ ID NO:5 consists of
20 8-15 amino acids. In some embodiments, the monomer which comprises SEQ ID NO:5 consists of 10-12 amino acid residues.

In preferred embodiments of the invention which relate to dimers, the monomer that comprises SEQ ID NO:5 comprises SEQ ID NO:4 or a fragment thereof such as one that has amino acids
25 7-12, 7-13, 7-14, 7-15, 7-16, 7-17, 7-18, 7-19, 6-12, 6-13, 6-14, 6-15, 6-16, 6-17, 6-18, 6-19, 5-12, 5-13, 5-14, 5-15, 5-16, 5-17, 5-18, 5-19, 4-12, 4-13, 4-14, 4-15, 4-16, 4-17, 4-18, 4-19, 3-12, 3-13, 3-14, 3-15, 3-16, 3-17, 3-18, 3-19, 2-12, 2-13, 2-14, 2-15, 2-16, 2-17, 2-18, 2-19, 1-12, 1-13, 1-14, 1-15, 1-
30 16, 1-17, 1-18 and 1-19 of SEQ ID NO:4.

In preferred embodiments of the invention which relate to dimers, the monomer that comprises at least a fragment of SEQ ID NO:3 consists of 4-25 amino acids and the monomer that comprises SEQ ID NO:5 consists of 6-25 amino acids. In some
35 embodiments, the monomer that comprises at least a fragment of SEQ ID NO:3 of at least 4 amino acids consists of 6-20 amino acids and the monomer which comprises SEQ ID NO:5 consists of

- 13 -

6-20 amino acids. In some embodiments, the monomer that comprises at least a fragment of SEQ ID NO:3 of at least 4 amino acids consists of 8-15 amino acids and the monomer that comprises SEQ ID NO:5 consists of 8-15 amino acids. In some
5 embodiments, the monomer that comprises at least a fragment of SEQ ID NO:3 of at least 4 amino acids consists of 10-12 amino acid residues and the monomer which comprises SEQ ID NO:5 consists of 10-12 amino acid residues.

In some preferred embodiments of the invention which
10 relate to dimers, the monomer that comprises at least a fragment of SEQ ID NO:3 of at least 4 amino acids comprises SEQ ID NO:3 or a fragment thereof such as one that has amino acids 1-4, 2-5, 3-6, 4-7, 5-8, 6-9, 7-10, 8-11, 1-5, 2-6, 3-7, 4-8, 5-9, 6-10, 7-11, 1-6, 2-7, 3-8, 4-9, 5-10, 6-11, 1-7, 2-8, 3-9,
15 4-10, 5-11, 1-8, 2-9, 3-10, 4-11, 1-9, 2-10, 3-11, 1-10 and 2-11 of SEQ ID NO:3 and the monomer that comprises SEQ ID NO:5 comprises SEQ ID NO:4 or a fragment thereof such as one that has amino acids 7-12, 7-13, 7-14, 7-15, 7-16, 7-17, 7-18, 7-19, 6-12, 6-13, 6-14, 6-15, 6-16, 6-17, 6-18, 6-19, 5-12, 5-13, 5-
20 14, 5-15, 5-16, 5-17, 5-18, 5-19, 4-12, 4-13, 4-14, 4-15, 4-16, 4-17, 4-18, 4-19, 3-12, 3-13, 3-14, 3-15, 3-16, 3-17, 3-18, 3-19, 2-12, 2-13, 2-14, 2-15, 2-16, 2-17, 2-18, 2-19, 1-12, 1-13, 1-14, 1-15, 1-16, 1-17, 1-18 and 1-19 of SEQ ID NO:4.

According to some embodiments of the invention,
25 peptides that inhibit IgE from binding to FcεR1 are provided which comprise at least a fragment of SEQ ID NO:3 of at least 4 amino acids and which further comprises SEQ ID NO:5. More particularly, in some embodiments of the invention, peptides comprise 10-100 amino acids including at least a fragment of
30 SEQ ID NO:3 of at least 4 amino acids and SEQ ID NO:5.

In some preferred embodiments of the invention, the peptides, which include at least a fragment of SEQ ID NO:3 of at least 4 amino acids and SEQ ID NO:5, comprise 10-50 amino acids including at least a fragment of SEQ ID NO:3 having 4-11
35 amino acids and at least a fragment of SEQ ID NO:4 having 6-19 amino acids including SEQ ID NO:5. In some embodiments, the peptides, which include at least a fragment of SEQ ID NO:3 of

- 14 -

at least 4 amino acids and SEQ ID NO:5, comprise 12-40 amino acids including at least a fragment of SEQ ID NO:1 of 6-20 amino acids including 6-11 amino acids from SEQ ID NO:3. In some embodiments, the peptide, which include at least a fragment of SEQ ID NO:3 of at least 4 amino acids and SEQ ID NO:5, comprise 14-30 amino acids including at least a fragment of SEQ ID NO:1 of 8-15 amino acids including 8-11 amino acids from SEQ ID NO:3. In some embodiments, the peptide, which include at least a fragment of SEQ ID NO:3 of at least 4 amino acids and SEQ ID NO:5, comprises 16-25 amino acids including at least a fragment of SEQ ID NO:1 of 11-12 amino acids including SEQ ID NO:3. In some embodiments, the peptide, which include at least a fragment of SEQ ID NO:3 of at least 4 amino acids and SEQ ID NO:5, comprises 12-40 amino acids including SEQ ID NO:5. In some embodiments, the peptide, which include at least a fragment of SEQ ID NO:3 of at least 4 amino acids and SEQ ID NO:5, comprises 14-30 amino acids including SEQ ID NO:5. In some embodiments, the peptide, which include at least a fragment of SEQ ID NO:3 of at least 4 amino acids and SEQ ID NO:5, comprises 16-25 amino acids including SEQ ID NO:5. In some embodiments, the peptide, which include at least a fragment of SEQ ID NO:3 of at least 4 amino acids and SEQ ID NO:5, comprises SEQ ID NO:3 or a fragment thereof such as one that has amino acids 1-4, 2-5, 3-6, 4-7, 5-8, 6-9, 7-10, 8-11, 1-5, 2-6, 3-7, 4-8, 5-9, 6-10, 7-11, 1-6, 2-7, 3-8, 4-9, 5-10, 6-11, 1-7, 2-8, 3-9, 4-10, 5-11, 1-8, 2-9, 3-10, 4-11, 1-9, 2-10, 3-11, 1-10 and 2-11 of SEQ ID NO:3 linked to SEQ ID NO:5 comprises SEQ ID NO:4 or a fragment thereof such as one that has amino acids 7-12, 7-13, 7-14, 7-15, 7-16, 7-17, 7-18, 7-19, 6-12, 6-13, 6-14, 6-15, 6-16, 6-17, 6-18, 6-19, 5-12, 5-13, 5-14, 5-15, 5-16, 5-17, 5-18, 5-19, 4-12, 4-13, 4-14, 4-15, 4-16, 4-17, 4-18, 4-19, 3-12, 3-13, 3-14, 3-15, 3-16, 3-17, 3-18, 3-19, 2-12, 2-13, 2-14, 2-15, 2-16, 2-17, 2-18, 2-19, 1-12, 1-13, 1-14, 1-15, 1-16, 1-17, 1-18 and 1-19 of SEQ ID NO:4.

In some embodiments, the peptide which includes the sequence which is defined as at least a fragment of SEQ ID NO:3 of at least 4 amino acids at the amino terminal portion of the

- 15 -

sequence relative to the position of the portion of the sequence that comprises SEQ ID NO:5.

In some embodiments, the peptide which includes the sequence SEQ ID NO:5 at the amino terminal portion of the sequence relative to the position of the portion of the sequence that which is defined as the sequence that comprises at least a fragment of SEQ ID NO:3 of at least 4 amino acids.

Peptides of the invention inhibit IgE from binding to FcεR1. The ability of a peptide to inhibit IgE from binding to FcεR1 may be determined by one having ordinary skill in the art by performing binding assays which are *in vitro* and/or *in vivo* assays using readily available starting materials. For example, assays to determine whether a peptide has biological activity are described in Examples 1 and 2. Briefly, an *in vitro* assay is described which allows one to determine whether or not IgE is inhibited from binding to FcεR1. An *in vivo* assay is described which allows one to determine whether or not GVHD occurs in a bone marrow transplant model. GVHD is linked to IgE dysregulation and to mast cell activity (Claman and Spiegelberg 1990 *Clin. Imm. Immunopath.* 56:46-53; and , Murphy et al. 1994 *J. Invest. Derma.* 102:451-461). Accordingly, inhibition of IgE-FcεR1 binding *in vivo* will prevent the lethal effects of GVHD in the *in vivo* assay. The ability of a compound of the invention to inhibit IgE from binding to FcεR1 refers its possession of IgE-FcεR1 binding inhibition activity in an IgE-FcεR1 binding assay.

In some embodiments, peptides of the present invention comprise D amino acids. As used herein, the term "D amino acid peptides" is meant to refer to peptides according to the present invention which comprise at least one and in some embodiments a plurality of D amino acids. D amino acid peptides inhibit IgE from binding to FcεR1, thus retaining the biological activity of peptides having L amino acid sequences. The use of D amino acid peptides is desirable as they are less vulnerable to degradation and therefore have a longer half life. D amino acid peptides comprising mostly all or consisting of D amino acids may comprise amino acid sequences

- 16 -

in the reverse order of SEQ ID NO:3 or fragments or derivatives thereof or SEQ ID NO:4 or fragments thereof.

In some embodiments, peptides of the present invention are conformationally restricted to present and maintain the proper structural conformation for biological activity. The compositions may comprise additional amino acid residues required to achieve proper three dimensional conformation including residues which facilitate circularization or desired folding. Accordingly, in preferred embodiments, cysteine residues are added to the terminal residues of peptides comprising SEQ ID NO:3 or fragments thereof or SEQ ID NO:4 of fragments thereof including SEQ ID NO:5. Formation of disulfide bonds between the terminal cysteines results in a circularized, i.e. cyclic peptides.

D amino acid residues may be provided to facilitate conformational restriction. For example, in order to facilitate disulfide bond formation and stability, a D amino acid cysteine may be provided at one or both termini.

Peptides which fall within the structural definition provided herein may be produced by those having ordinary skill in the art using routine techniques and readily available starting materials without undue experimentation. One having ordinary skill in the art can then test the peptides to determine whether or not they inhibit IgE binding to FcεR1 and are thus peptides of the invention.

It is also contemplated that some embodiments of the invention include the peptides that comprise at least a fragment of a derivative of SEQ ID NO:3 of at least 4 amino acids rather than a fragment of SEQ ID NO:3 and/or a derivative of SEQ ID NO:5 rather than SEQ ID NO:5. As used herein, the term "derivative of SEQ ID NO:3" is meant to refer to a peptide which comprises an amino acid sequence substantially identical to SEQ ID NO:3. As used herein, the term "derivative of SEQ ID NO:5" is meant to refer to a peptide which comprises an amino acid sequence substantially identical to SEQ ID NO:5. As used herein, the term "substantially identical to SEQ ID NO:3 or a fragment thereof" refers to an amino acid sequence that is the

- 17 -

same as the amino acid sequence of SEQ ID NO:3 or a fragment thereof except some of the residues are deleted, substituted with conservative amino acids or additional amino acids are inserted. As used herein, the term "substantially identical to

5 SEQ ID NO:5" refers to an amino acid sequence that is the same as the amino acid sequence of SEQ ID NO:5 except some of the residues are deleted, substituted with conservative amino acids or additional amino acids are inserted. Those having ordinary skill in the art can readily design and produce derivatives

10 having substantially identical amino acid sequences of SEQ ID NO:3 or a fragment thereof or SEQ ID NO:5 with deletions and/or insertions and/or conservative substitutions of amino acids. For example, following what are referred to as Dayhof's rules for amino acid substitution (Dayhof, M.D. (1978) *Nat. Biomed.*

15 *Res. Found.*, Washington, D.C. Vol. 5, supp. 3), amino acid residues in a peptide sequence may be substituted with comparable amino acid residues. Such substitutions are well known and are based the upon charge and structural characteristics of each amino acid. Derivatives include

20 fragments of SEQ ID NO:3 or SEQ ID NO:5 with deletions and/or insertions and/or conservative substitutions.

Peptides may be synthesized by well known techniques such as the solid-phase synthetic technique initially described by Merrifield, in *J. Am. Chem. Soc.*, 15:2149-2154 (1963).

25 Other peptide synthesis techniques may be found, for example, in M. Bodanszky et al., (1976) *Peptide Synthesis*, John Wiley & Sons, 2d Ed.; Kent and Clark-Lewis in *Synthetic Peptides in Biology and Medicine*, p. 295-358, eds. Alitalo, K., et al. Science Publishers, (Amsterdam, 1985); as well as other

30 reference works known to those skilled in the art. A summary of peptide synthesis techniques may be found in J. Stuart and J.D. Young, *Solid Phase Peptide Synthesis*, Pierce Chemical Company, Rockford, IL (1984), which is incorporated herein by reference. The synthesis of peptides by solution methods may

35 also be used, as described in *The Proteins*, Vol. II, 3d Ed., p. 105-237, Neurath, H. et al., Eds., Academic Press, New York, NY (1976). Appropriate protective groups for use in such

- 18 -

syntheses will be found in the above texts, as well as in J.F.W. McOmie, *Protective Groups in Organic Chemistry*, Plenum Press, New York, NY (1973), which is incorporated herein by reference.

5 In general, these synthetic methods involve the sequential addition of one or more amino acid residues or suitable protected amino acid residues to a growing peptide chain. Normally, either the amino or carboxyl group of the first amino acid residue is protected by a suitable,
10 selectively removable protecting group. A different, selectively removable protecting group is utilized for amino acids containing a reactive side group, such as lysine.

 Using a solid phase synthesis as an example, the protected or derivatized amino acid is attached to an inert
15 solid support through its unprotected carboxyl or amino group. The protecting group of the amino or carboxyl group is then selectively removed and the next amino acid in the sequence having the complementary (amino or carboxyl) group suitably protected is admixed and reacted with the residue already
20 attached to the solid support. The protecting group of the amino or carboxyl group is then removed from this newly added amino acid residue, and the next amino acid (suitably protected) is then added, and so forth. After all the desired amino acids have been linked in the proper sequence, any
25 remaining terminal and side group protecting groups (and solid support) are removed sequentially or concurrently, to provide the final peptide. The peptide of the invention is preferably devoid of benzylated or methylbenzylated amino acids. Such protecting group moieties may be used in the course of
30 synthesis, but they are removed before the peptides are used. Additional reactions may be necessary, as described elsewhere, to form intramolecular linkages to restrain conformation.

 Peptides may also be prepared by recombinant DNA techniques. Provision of a suitable DNA sequence encoding the
35 desired peptide permits the production of the peptide using recombinant techniques now known in the art. The coding sequence can be obtained from natural sources or synthesized or

- 19 -

otherwise constructed using widely available starting materials by routine methods. When the coding DNA is prepared synthetically, advantage can be taken of known codon preferences of the intended host where the DNA is to be expressed.

To produce peptides of the invention, one having ordinary skill in the art can, using well known techniques, synthesize or obtain a DNA molecule encoding SEQ ID NO:1 or portions thereof which include SEQ ID NO:3 or a fragment thereof or SEQ ID NO:5 from readily available human DNA and insert that DNA molecule into a commercially available expression vector for use in well known expression systems. Including in these systems are those in which the peptide of interest is produced as a fusion protein.

For example, the commercially available plasmid pSE420 (Invitrogen, San Diego, CA) may be used for recombinant production in *E. coli*. The commercially available plasmid pYES2 (Invitrogen, San Diego, CA) may be used for production in *S. cerevisiae* strains of yeast. The commercially available MaxBac™ (Invitrogen, San Diego, CA) complete baculovirus expression system may be used for production in insect cells. The commercially available plasmid pcDNA I (Invitrogen, San Diego, CA) may be used for production in mammalian cells such as Chinese Hamster Ovary cells.

One having ordinary skill in the art may use these or other commercially available expression vectors and systems or produce vectors using well known methods and readily available starting materials. Expression systems containing the requisite control sequences, such as promoters and polyadenylation signals, and preferably enhancers, are readily available and known in the art for a variety of hosts. See e.g., Sambrook et al., *Molecular Cloning a Laboratory Manual*, Second Ed. Cold Spring Harbor Press (1989). Thus, the desired proteins can be prepared in both prokaryotic and eukaryotic systems, resulting in a spectrum of processed forms of the protein.

- 20 -

The most commonly used prokaryotic system remains *E. coli*, although other systems such as *B. subtilis* and *Pseudomonas* are also useful. Suitable control sequences for prokaryotic systems include both constitutive and inducible
5 promoters including the *lac* promoter, the *trp* promoter, hybrid promoters such as *tac* promoter, the *lambda* phage P1 promoter. In general, foreign proteins may be produced in these hosts either as fusion or mature proteins. When the desired sequences are produced as mature proteins, the sequence
10 produced may be preceded by a methionine which is not necessarily efficiently removed. Accordingly, the peptides and proteins claimed herein may be preceded by an N-terminal Met when produced in bacteria. Moreover, constructs may be made wherein the coding sequence for the peptide is preceded by an
15 operable signal peptide which results in the secretion of the protein. When produced in prokaryotic hosts in this matter, the signal sequence is removed upon secretion.

A wide variety of eukaryotic hosts are also now available for production of recombinant foreign proteins. As
20 in bacteria, eukaryotic hosts may be transformed with expression systems which produce the desired protein directly, but more commonly signal sequences are provided to effect the secretion of the protein. Eukaryotic systems have the additional advantage that they are able to process introns
25 which may occur in the genomic sequences encoding proteins of higher organisms. Eukaryotic systems also provide a variety of processing mechanisms which result in, for example, glycosylation, carboxy-terminal amidation, oxidation or derivatization of certain amino acid residues, conformational
30 control, and so forth.

Commonly used eukaryotic systems include, but is not limited to, yeast, fungal cells, insect cells, mammalian cells, avian cells, and cells of higher plants. Suitable promoters are available which are compatible and operable for use in each
35 of these host types as well as are termination sequences and enhancers, as e.g. the baculovirus polyhedron promoter. As above, promoters can be either constitutive or inducible. For

- 21 -

example, in mammalian systems, the mouse metallothionein promoter can be induced by the addition of heavy metal ions.

In some embodiments, a DNA molecule that includes a nucleotide sequence which encodes the peptide of the invention is synthesized using the amino acid sequence information herein and the genetic code. Those having ordinary skill in the art can readily synthesize DNA molecules that include nucleotide sequences which encode the peptides of the invention using codons preferred by a desired host cell. The DNA molecule that is generated may be inserted in an expression vector which allows for very high levels of expression, sometimes referred to as overexpression, in a desired host.

The particulars for the construction of expression systems suitable for desired hosts are known to those in the art. For recombinant production of the protein, the DNA encoding it is suitably ligated into the expression vector of choice and then used to transform the compatible host which is then cultured and maintained under conditions wherein expression of the foreign gene takes place. The protein of the present invention thus produced is recovered from the culture, either by lysing the cells or from the culture medium as appropriate and known to those in the art.

One having ordinary skill in the art can, using well known techniques, isolate the protein that is produced.

In addition to peptides, the present invention contemplates compounds which display substantially the same surface as the peptides of the invention. Such compounds, referred to as mimetics, are not peptides but comprise a similar surface as the peptides and can thus interact with other molecules in a similar fashion as the peptides of the invention. By providing a similar surface involved in intermolecular interactions, mimetics perform essentially the same function by essentially the same means to achieve essentially the same result as the peptides of the invention.

The present invention relates to a method of therapeutically or prophylactically treating an individual suffering from or susceptible to an immediate allergic

- 22 -

response. Those having ordinary skill in the art can readily identify individuals suspected of suffering from or being susceptible to an immediate allergic response conditions such as response is selected from the group consisting of: graft
5 versus host disease, allergic rhinitis, food allergies, atropic dermatitis and allergic asthma. In the case of GVHD, treatment may be provided prophylactically in conjunction with transplantation procedure or in response to symptoms associated with GVHD. Those with ordinary skill in the art could readily
10 identify individuals for whom administration of the peptides of the invention would be beneficial to alleviate or prevent immediate allergy response conditions.

The method of therapeutically or prophylactically treating an individual suffering from or susceptible to an
15 immediate allergic response comprises administering to said individual an effective amount of a peptide according to the invention. A prophylactically effective amount is an amount which is effective to prevent or decrease the immediate allergic response in an individual susceptible to such a
20 reaction without causing lethal side effects on the individual. A prophylactically effective amount is an amount which is effective to decrease or eliminate the immediate allergic response in an individual suffering from such a reaction without causing lethal side effects on the individual. Those
25 having ordinary skill in the art can readily and routinely determine the ranges of both prophylactically and therapeutically effective amounts of the peptides of the invention without undue experimentation.

It is contemplated that the present invention is
30 useful to therapeutically or prophylactically treat individuals suspected of suffering from or being susceptible to reactions are mediated through the interaction of immunoglobulin E (IgE) with the α -chain of its high affinity Fc receptor, Fc ϵ R1.

The present invention provides pharmaceutical
35 compositions that comprise the peptides of the invention and pharmaceutically acceptable carriers or diluents. The pharmaceutical composition of the present invention may be

- 23 -

formulated by one having ordinary skill in the art. Suitable pharmaceutical carriers are described in *Remington's Pharmaceutical Sciences*, A. Osol, a standard reference text in this field, which is incorporated herein by reference. In
5 carrying out methods of the present invention, conjugated compounds of the present invention can be used alone or in combination with other diagnostic, therapeutic or additional agents. Such additional agents include excipients such as coloring, stabilizing agents, osmotic agents and antibacterial
10 agents.

For parenteral administration, the peptides of the invention can be, for example, formulated as a solution, suspension, emulsion or lyophilized powder in association with a pharmaceutically acceptable parenteral vehicle. Examples of
15 such vehicles are water, saline, Ringer's solution, dextrose solution, and 5% human serum albumin. Liposomes and nonaqueous vehicles such as fixed oils may also be used. The vehicle or lyophilized powder may contain additives that maintain isotonicity (e.g., sodium chloride, mannitol) and chemical
20 stability (e.g., buffers and preservatives). The formulation is sterilized by commonly used techniques. For example, a parenteral composition suitable for administration by injection is prepared by dissolving 1.5% by weight of active ingredient in 0.9% sodium chloride solution.

25 The pharmaceutical compositions according to the present invention may be administered as a single dose or in multiple doses. The pharmaceutical compositions of the present invention may be administered either as individual therapeutic agents or in combination with other therapeutic agents. The
30 treatments of the present invention may be combined with conventional therapies, which may be administered sequentially or simultaneously.

The pharmaceutical compositions of the present invention may be administered by any means that enables the
35 active agent to reach the targeted cells. These methods include, but are not limited to, oral, topical, intradermal, subcutaneous, intravenous, intramuscular and intraparenteral

- 24 -

modes of administration. The compounds may be administered singly or in combination with other compounds. The compounds of the invention are preferably administered with a pharmaceutically acceptable carrier selected on the basis of the selected route of administration and standard pharmaceutical practice.

The dosage administered varies depending upon factors such as: pharmacodynamic characteristics; its mode and route of administration; age, health, and weight of the recipient; nature and extent of symptoms; kind of concurrent treatment; and frequency of treatment. Usually, the dosage of peptide can be about 1 to 3000 milligrams per 50 kilograms of body weight; preferably 10 to 1000 milligrams per 50 kilograms of body weight; more preferably 25 to 800 milligrams per 50 kilograms of body weight. Ordinarily 8 to 800 milligrams are administered to an individual per day in divided doses 1 to 6 times a day or in sustained release form is effective to obtain desired results.

Examples

20 Example 1

In vitro inhibition of binding of IgE Fc to Fc ϵ by biosensor assay.

The kinetics of IgE Fc binding to soluble Fc ϵ were determined using a BIAcore biosensor (Pharmacia Biosensor AB, Uppsala, Sweden). The Fc ϵ RI α was immobilized via primary amine groups to a carboxymethylated dextran-coated CM5 sensor chip, at a concentration of 80 μ g/ml in 10mM sodium acetate buffer pH 4.5, using N-hydroxy-succinimide and N-ethyl-N'-(3-diethylaminopropyl)-carbodiimide (Pharmacia Amine Coupling Kit). Free sites were blocked with 1M ethanolamine-HCl pH 8.5. Binding of peptides to IgE was measured by the ability of peptides to prevent binding of IgE to the immobilized Fc ϵ RI α . Various concentrations of peptide were incubated with a fixed concentration of IgE (10nM) and allowed to equilibrate for 24h. The peptide-IgE mixtures were injected over the sensor surface, in 10mM HEPES buffered saline (HBS) with 150mM NaCl, 3.4 mM

- 25 -

EDTA and 0.005% surfactant P20 (Pharmacia), at 25°C, using conditions under which binding is limited by mass transport rather than by kinetic parameters. This allowed measurement of the concentration of free IgE, and thus the solution
5 association constant (K_a) for the receptor-ligand interaction. The initial rate of IgE binding to the immobilized receptor is presented as a function of peptide concentration in Figure 4.

Example 2 In vivo test to identify IgE-IgE Receptor Binding Inhibitors

10 The transplantation of B10.D2 CD4+ T-cells along with T-cell-depleted bone marrow has been shown previously to induce a lethal GVHD response in irradiated (850 cGy) DBA/2 mice (Korngold and Sprent (1987) *J. Exp. Med.* 165:1552-1564, which is incorporated herein by reference). In addition, in this
15 strain combination dermal mast cells are believed to be involved in the early stages of cutaneous GVHD development. The transplantation of 7×10^6 B10.D2 CD4+ T-cells resulted in a 100% incidence of lethal GVHD in DBA/2 recipients with a median survival time of 9 days. Transplantation of donor T-
20 cell-depleted bone marrow alone resulted in complete survival out to termination of the experiment at 70 days. To determine the ability of an L form and a reverse D form of a test compound of the invention to inhibit IgE-IgE receptor binding and thereby inhibit GVHD, DBA/2 recipients of B10.D2 CD4+ T-
25 cells are treated on days 0 and 3 with 0.5 mg of either test compound i.v. Incidence of survival and median survival time are compared to an untreated group and a group treated with a scrambled reverse D form of the test compound. These data indicate that the ability of a reverse D test compound to
30 effect the development of GVHD and alloreactivity of transplanted donor T-cells.

Example 3

 In one embodiment of the invention, a peptide that includes SEQ ID NO:3 and that is restrained by cysteines at
35 each end is synthesized. The peptide and has the following sequence:

- 26 -

C-I-Y-Y-K-D-G-E-A-L-K-Y-C^D, where the carboxy terminal cysteine is a "D" amino acid. This D amino acid peptide is referred to herein as D amino acid peptide #1. D amino acid peptide #1 consists of a total of 13 amino acids and comprises all of SEQ ID NO:3. D amino acid peptide #1 is conformationally restricted using two terminal cysteines to form a disulfide bond which cyclicizes the peptide. The D amino acid cysteine is included to facilitate the formation and stability of the disulfide bond.

10 Approximately 0.5×10^6 transfected CHO cells were incubated with a mixture of 5 nM labelled myeloma IgE and D amino acid peptide #1 was titrated at various concentrations for 1h at room temperature. Non-specific bindings were determined by pre-incubating the cells with a 100-fold excess
15 of rat IgE. After incubation, the cells were separated from the unbound iodinated IgE by 2 min. centrifugation through mixed oils. The cell pellets were then counted in a 5500 Beckman gamma counter.

Concentration of the D amino acid peptide #1 used
20 ranged from 9.324×10^{-8} M to 3.73×10^{-5} M. The $IC_{50} = 5.14 \times 10^{-6}$ M. The results are shown on Figure 1.

As shown in Figure 2, the transplantation of 2×10^6 B10.D2 CD4+ T cells resulted in an 83% incidence of lethal GVHD in DBA/2 recipients with a median survival time of 15 days.
25 Transplantation of donor T-cell-depleted bone marrow alone resulted in complete survival out to termination of the experiment at 80 days. DBA/2 recipients of B10.D2 CD4+ T cells and treated on days 0 and 3 with 0.5 mg D amino acid peptide #1 i.v. exhibited a significantly higher incidence of survival
30 (67%) and a significantly prolonged median survival time of greater than 80 days ($p=0/04$), compared to the untreated group. This data indicates that the IgE receptor analogs of the invention can diminish the development of GVHD and alloreactivity of transplanted donor T cells.

35 Example 4

Surface plasmon resonance experiment

- 27 -

The kinetics of binding of IgE Fc to soluble FcεR1α were determined using a BIAcore biosensor (Pharmacia Biosensor AB, Uppsala, Sweden). The FcεR1α was immobilized via primary amine groups to a carboxymethylated dextran-coated CM5 sensor chip, at a concentration of 85 μg/ml in 10mM sodium acetate buffer pH 4.5, using N-hydroxy-succinimide and N-ethyl-N'-(3-diethyl-aminopropyl)-carbodiimide. Free sites were blocked with 1M ethanolamine-hydrochloride pH 8.5.

The association of IgE-Fc at a concentration of 5nM was monitored for 6 minutes in the presence and absence of a derivative of the D amino acid peptide #1 at various concentrations (at 24°C). The derivative of D amino acid peptide #1 is identical to D amino acid #1, which is described in Example 3, except it is biotinylated. At a concentration of 2.5mM, virtually complete inhibition of IgE-Fc binding was observed. The results are shown in Figure 3.

Example 5

SEQ ID NO:6 is SEQ ID NO:3 with cysteine residues at the N and C terminals. The two cysteines form disulfide bonds which result in the cyclization of the peptide.

D amino acid peptide #1 has the same amino acid sequence as SEQ ID NO:6 except that the C terminus cysteine is a D amino acid. D amino acid peptide #1 is also referred to as cyclo(L-262).

D amino acid peptide #2 has the sequence of D amino acids from amino terminus to carboxy terminus: Cys Tyr Lys Leu Ala Glu Gly Asp Lys Tyr Tyr Ile Cys_(L). The C terminus cysteine is an L amino acid. This sequence, also referred to as cyclo(rD-262), is the reverse amino acid sequence of SEQ ID NO:6.

Peptides cyclo(L-262) and cyclo(rD-262) were synthesized and tested for activity. The sequences of the peptides are as follows, cyclo(L-262) CIYYKDGEALKYC_(D) (all L amino acids except the carboxy terminal cysteine) and cyclo(rD-262) CYKLAEGDKYYIC_(L) (all D amino acids except carboxy terminal cysteine).

- 28 -

The following scrambled peptides were used as controls: cyclo(rD-scram) CLEADYKGYKYIC_(L) (all D amino acids except carboxy terminal cysteine) and cyclo(L-scram) CIYKYGKYDAELC_(D) (all L amino acids except the carboxy terminal 5 cysteine).

Peptides were synthesized on an Applied Biosystems 430A fully automated peptide synthesizer using standard fmoc chemistry. Intramolecular disulfides were enriched by an air oxidation refolding procedure carried out at 100 µg/ml in 50 mM ammonium bicarbonate, pH 8.5, stirred overnight at 23°C. Peptides show greater than 95% intramolecular disulfide bonding at the end of this procedure as monitored by Ellman's reagent and HPLC analysis. In addition, matrix assisted laser desorption/ionization mass spectrometry (MALDI-MS) was used to confirm the identity and cyclized status of the peptide product. The intramolecular disulfide cyclized product was then purified by RP-HPLC.

Example 6

In vivo inhibition of GVHD (Figure 5).

Bone marrow cells were obtained from the femurs and tibiae of donor mice by flushing with buffered saline solution (BSS) supplemented with 0.1% bovine serum albumin (Hyclone, Logan, UT). To prepare anti-Thy-1 (T cell-depleted) bone-marrow (ATBM), cells were incubated with J1J mAb (diluted 1:100) and complement 1:25 for 45 minutes at 37°C, followed by four washes. CD4+ T cell-enriched donor cell populations were prepared by initially treating pooled spleen and lymph node cells with the anti-B cell J1ld mAb (diluted 1:5) and complement (1:25) for 1 hour at 37°C. This treatment generally resulted in a population between 90-95% Thy-1+ cells, as measured for phenotyping by flow cytometric analysis. These cells were further treated twice with anti-CD8 mAb (3.268, diluted 1:50) plus complement (1:25) for 45 minutes at 37°C, initially and again for 30 minutes with a wash in between. The enriched CD4+ T cells were phenotyped by flow cytometry analysis and were routinely found to be negative for CD8+ T cell subset. For GVHD induction, DBA/2 recipient mice were

- 29 -

irradiated 900 cGy by exposure from a Gammacell ^{137}Cs source (130 cGy/min). Approximately 6 hours later these mice were injected intravenously with donor B10.D2 ATBM cells (2×10^6) alone as a negative control, or a mixture of ATBM (2×10^6) together with 5×10^6 CD4+ T cells.

To determine the ability of cyclo(L-262) and cyclo(rD-262) peptides, and scrambled versions of these peptides, to inhibit receptor binding and thereby inhibit GVHD, DBA/2 recipients of B10.D2 CD4+ T-cells were treated on days 0 and 3 with 0.5 mg of peptide i.v. Mice were observed daily for mortality until 70 days post-transplantation and weights were monitored twice weekly. Data are shown in Figure 5.

Example 7

In vivo inhibition of antigen-induced anaphylaxis (Figure 6). BALB/c mice primed to bacteriophage lambda cI repressor peptide 9-29 (ref. J. Exp. Med., 174, 847, 1991), with 200 μg i.p., followed by a secondary boost 3 weeks later. They were subsequently challenged with 250 μg of the same antigen, and scored for level of distress (level 4 = severe fatal reaction) for up to 40 minutes after challenge. Cyclo(rD-262) peptide (0.5 mg) was administered at either time 0 or 15 minutes before challenge, and compared with administration of a scrambled version of the peptide at time 0. The results are shown in Figure 6.

- 30 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Jameson, Bradford A.
Sutton, Brian J.
McDonnell, James M.
Gould, Hannah J.
Korngold, Robert
Beavil, Andrew J.

(ii) TITLE OF INVENTION: IgE Antagonists

(iii) NUMBER OF SEQUENCES: 6

(iv) CORRESPONDENCE ADDRESS:
(A) ADDRESSEE: Woodcock Washburn Kurtz Mackiewicz & Norris
(B) STREET: One Liberty Place, 46th Floor
(C) CITY: Philadelphia
(D) STATE: PA
(E) COUNTRY: USA
(F) ZIP: 19103

(v) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: Wordperfect

(vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: US 08/271,943
(B) FILING DATE: 08-JUL-1994
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:
(A) NAME: DeLuca, Mark
(B) REGISTRATION NUMBER: 33,229
(C) REFERENCE/DOCKET NUMBER: TJU-1597

(ix) TELECOMMUNICATION INFORMATION:
(A) TELEPHONE: 215-568-3100
(B) TELEFAX: 215-568-3439

(2) INFORMATION FOR SEQ ID NO:1:

```

40      (i) SEQUENCE CHARACTERISTICS:
          (A) LENGTH: 193 amino acids
          (B) TYPE: amino acid
          (D) TOPOLOGY: linear
      (ii) MOLECULE TYPE: protein
      (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

45      Glu Ser Pro Thr Leu Leu Cys Val Ala Leu Leu Phe Phe Ala Pro Asp
          1             5             10             15

          Gly Val Leu Ala Val Pro Gln Lys Pro Lys Val Ser Leu Asn Pro Pro
              20             25             30

          Trp Asn Arg Ile Phe Lys Gly Glu Asn Val Thr Leu Thr Cys Asn Gly
              35             40             45

50      Asn Asn Phe Phe Glu Val Ser Ser Thr Lys Trp Phe His Asn Gly Ser
          50             55             60

          Leu Ser Glu Glu Thr Asn Ser Ser Leu Asn Ile Val Met Ala Lys Phe
          65             70             75             80

55      Glu Asp Ser Gly Glu Tyr Lys Cys Gln His Gln Gln Val Asn Glu Ser
              85             90             95

          Glu Pro Val Tyr Leu Glu Val Phe Ser Asp Trp Leu Leu Leu Gln Ala
              100            105            110

```


- 31 -

Ser Ala Glu Val Val Met Glu Gly Gln Pro Leu Phe Leu Arg Cys His
 115 120 125

Gly Trp Arg Met Trp Asp Val Tyr Lys Val Ile Tyr Tyr Lys Asp Gly
 130 135 140

5 Glu Ala Leu Lys Tyr Trp Tyr Glu Asn His Asn Ile Ser Ile Thr Asn
 145 150 155 160

Ala Thr Val Glu Asp Ser Gly Thr Tyr Tyr Cys Thr Gly Lys Val Trp
 165 170 175

10 Gln Leu Asp Tyr Glu Ser Glu Pro Leu Asn Ile Thr Val Ile Lys Ala
 180 185 190

Pro

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 174 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

20 Asp Ser Gly Thr Val Trp Gly Ala Leu Gly His Gly Ile Asn Leu Asn
 1 5 10 15

Ile Pro Asn Gly Gln Met Thr Asp Asp Ile Asp Glu Val Arg Trp Glu
 20 25 30

25 Arg Gly Ser Thr Leu Val Ala Glu Lys Arg Lys Met Lys Pro Phe
 35 40 45

Leu Lys Ser Gly Ala Phe Glu Ile Leu Ala Asn Gly Asp Leu Lys Ile
 50 55 60

Lys Asn Leu Thr Arg Asp Asp Ser Gly Thr Tyr Asn Val Thr Val Tyr
 65 70 75 80

30 Ser Thr Asn Gly Thr Arg Ile Leu Asn Lys Ala Leu Asp Leu Arg Ile
 85 90 95

Leu Glu Met Val Ser Lys Pro Met Ile Tyr Trp Glu Cys Ser Asn Ala
 100 105 110

35 Thr Leu Thr Cys Glu Val Leu Glu Gly Thr Asp Val Glu Leu Lys Leu
 115 120 125

Tyr Gln Gly Lys Glu His Leu Arg Ser Leu Arg Gln Lys Thr Met Ser
 130 135 140

Tyr Gln Trp Thr Asn Leu Arg Ala Pro Phe Lys Cys Lys Ala Val Asn
 145 150 155 160

40 Arg Val Ser Gln Glu Ser Glu Met Glu Val Val Asn Cys Pro
 165 170

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- 45 (A) LENGTH: 11 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

- 32 -

Ile Tyr Tyr Lys Asp Gly Glu Ala Leu Lys Tyr
 1 5 10

(2) INFORMATION FOR SEQ ID NO:4:

- 5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

10 Thr Tyr Tyr Cys Thr Gly Lys Val Trp Gln Leu Asp Tyr Glu Ser Glu
 1 5 10 15

Pro Leu Asn

(2) INFORMATION FOR SEQ ID NO:5:

- 15 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 6 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

20 Lys Val Trp Gln Leu Asp
 1 5

(2) INFORMATION FOR SEQ ID NO:6:

- 25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 13 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

30 Cys Ile Tyr Tyr Lys Asp Gly Glu Ala Leu Lys Tyr Cys
 1 5 10

- 33 -

Claims

1. A peptide consisting of 4-50 amino acids, wherein said peptide
 - a) comprises at least a fragment of SEQ ID NO:3 of at least 4 amino acids and
 - b) is capable of modulating IgE binding to FcεR1.
2. The peptide of claim 1 wherein said peptide consists of 4-25 amino acids.
3. The peptide of claim 1 wherein said peptide consists of 8-15 amino acids.
4. The peptide of claim 1 wherein said peptide comprises SEQ ID NO:3 or a fragment thereof selected from the group consisting of amino acids 1-4, 2-5, 3-6, 4-7, 5-8, 6-9, 7-10, 8-11, 1-5, 2-6, 3-7, 4-8, 5-9, 6-10, 7-11, 1-6, 2-7, 3-8, 4-9, 5-10, 6-11, 1-7, 2-8, 3-9, 4-10, 5-11, 1-8, 2-9, 3-10, 4-11, 1-9, 2-10, 3-11, 1-10, 2-11 of SEQ ID NO:3..
5. The peptide of claim 1 wherein said peptide comprises SEQ ID NO:3.
6. The peptide of claim 1 wherein said peptide is conformationally restricted.
7. The peptide of claim 1 wherein said peptide has an amino terminal cysteine and a carboxy terminal cysteine which are linked by a disulfide bond.
8. The peptide of claim 1 having the amino acid sequence:
C-I-Y-Y-K-D-G-E-A-L-K-Y-C^D, wherein the carboxy terminal cysteine is a "D" amino acid.
9. The peptide of claim 1 wherein said peptide is a D amino acid having the amino acid sequence: C^L-Y-K-L-A-E-G-D-K-

- 34 -

Y-Y-I-C, wherein the amino terminal cysteine is an "L" amino acid.

10. A pharmaceutical composition comprising:

a) a peptide consisting of 4-50 amino acids, wherein
5 said peptide

i) comprises at least a fragment of SEQ ID NO:3
of at least 4 amino acids and

ii) is capable of modulating IgE binding to
FcεR1; and

10 b) a pharmaceutically acceptable carrier or diluent.

11. The pharmaceutical composition of claim 10 wherein
said peptide has the amino acid sequence: C-I-Y-Y-K-D-G-E-A-L-
K-Y-C^D, wherein the carboxy terminal cysteine is a "D" amino
acid.

15 12. The pharmaceutical composition of claim 10 wherein
said peptide is a D amino acid having the amino acid sequence:
C^L-Y-K-L-A-E-G-D-K-Y-Y-I-C, wherein the amino terminal cysteine
is an "L" amino acid.

13. A peptide consisting of 6-50 amino acids, wherein said
20 peptide

a) comprises SEQ ID NO:5 and

b) is capable of modulating IgE binding to FcεR1.

14. The peptide of claim 13 wherein said peptide consists
of 6-20 amino acids.

25 15. The peptide of claim 13 wherein said peptide consists
of 8-15 amino acids.

16. The peptide of claim 13 wherein said peptide comprises
SEQ ID NO:4 or a fragment thereof such as one that has amino
acids 7-12, 7-13, 7-14, 7-15, 7-16, 7-17, 7-18, 7-19, 6-12, 6-
30 13, 6-14, 6-15, 6-16, 6-17, 6-18, 6-19, 5-12, 5-13, 5-14, 5-15,

- 35 -

5-16, 5-17, 5-18, 5-19, 4-12, 4-13, 4-14, 4-15, 4-16, 4-17, 4-18, 4-19, 3-12, 3-13, 3-14, 3-15, 3-16, 3-17, 3-18, 3-19, 2-12, 2-13, 2-14, 2-15, 2-16, 2-17, 2-18, 2-19, 1-12, 1-13, 1-14, 1-15, 1-16, 1-17, 1-18 or 1-19 of SEQ ID NO:4.

5 17. The peptide of claim 13 wherein said peptide is conformationally restricted.

18. The peptide of claim 13 wherein said peptide has an amino terminal cysteine and a carboxy terminal cysteine which are linked by a disulfide bond.

10 19. A pharmaceutical composition comprising:
a) a peptide consisting of 6-50 amino acids, wherein said peptide

i) comprises SEQ ID NO:5 and

ii) is capable of modulating IgE binding to

15 FcεR1; and

b) a pharmaceutically acceptable carrier or diluent.

20. A method of inhibiting IgE binding to FcεR1 on cells comprising the steps of:

contacting cells that contain the FcεR1 with an amount
20 of a peptide sufficient to inhibiting IgE binding to FcεR1, wherein said peptide

a) consists of 4-50 amino acids,

b) comprises at least a fragment of SEQ ID NO:3 of at least 4 amino acids.

25 21. A method of inhibiting IgE binding to FcεR1 on cells comprising the steps of:

contacting cells that contain the FcεR1 with an amount of a peptide sufficient to inhibiting IgE binding to FcεR1, wherein said peptide

30 a) consists of 6-50 amino acids,

b) comprises SEQ ID NO:5.

- 36 -

22. A method of therapeutically or prophylactically treating an individual suspected of suffering from or being susceptible to an immediate allergic response comprising the steps of:

5 administering to said individual an effective amount of a peptide which:

a) consists of 4-50 amino acids,

b) comprises at least a fragment of SEQ ID NO:3 of at least 4 amino acids, and

10 c) inhibits IgE binding to FcεR1.

23. The method of claim 22 wherein said immediate allergic response is selected from the group consisting of: graft versus host disease, allergic rhinitis, food allergies, atropic dermatitis and allergic asthma.

15 24. A method of therapeutically or prophylactically treating an individual suspected of suffering from or being susceptible to an immediate allergic response comprising the steps of:

administering to said individual an effective amount
20 of a peptide which:

a) consists of between 6 and 50 amino acids,

b) comprises SEQ ID NO:5, and

c) inhibits IgE binding to FcεR1.

25. The method of claim 24 wherein said immediate allergic
25 response is selected from the group consisting of: graft versus host disease, allergic rhinitis, food allergies, atropic dermatitis and allergic asthma.

26. A peptide dimer comprising a first monomer and a second monomer wherein

30 a) said first monomer

i) consists of 4-50 amino acids,

ii) comprises at least a fragment of SEQ ID NO:3 of at least 4 amino acids and

- 37 -


b) said second monomer

i) consists of 6-50 amino acids,

ii) comprises SEQ ID NO:5;


wherein said dimer is capable of inhibiting IgE binding to
5 FcεR1.

1/6

***Fig. 1***

2/6


FIGURE 2



3/6

FIGURE 3

Name: Sensorgram 19: RUNNING




FcεRIα
immobilised
on sensor
surface

↑
In presence of
2.5 mM #262-86
2.5 nM IgE Fc


↑
In absence of
inhibitor.
5 nM IgE Fc

FIGURE 4

Inhibition of IgE-Fc binding to sFcεRIα
by receptor derived peptides




5/6

***Fig. 5***

6/6

FIGURE 6

Inhibition of antigen-induced anaphylaxis
by receptor-derived peptides



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/08401

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 38/00, 38/04; C07K 5/00, 7/00, 16/00

US CL :530/324, 326, 327, 328, 329, 330; 514/12, 13, 14, 15, 16, 17, 18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 530/324, 326, 327, 328, 329, 330; 514/12, 13, 14, 15, 16, 17, 18

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, STN

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US, A, 5,292,867 (CHANG) 08 MARCH 1994, see entire patent.	1-26

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

19 OCTOBER 1995

Date of mailing of the international search report

31 OCT 1995

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

BENET PRICKRIL

Telephone No. (703) 308-0196